ABRF-2003 ESRG: Analysis of a PVDF-Bound Known Protein with a Single Amino-Terminus.

Scott D. Buckel[1], Richard G. Cook[2], J. Myron Crawford[3], Nancy Denslow[4], Joseph Fernandez[5], Ben J. Madden[6], Laurey Steinke[7], and John M. Neveu[8]

[1]Amgen, Boulder, CO, [2]Baylor College of Medicine, Houston, TX, [3]Yale University, New Haven, CT, [4]University of Florida, Gainesville, FL, [5]Rockefeller University, New York, NY, [6]Mayo Clinic, Rochester, MN, [7]Nebraska Medical Center, Omaha, NE, [8]Harvard University, 16 Divinity Ave., Cambridge, MA 02138.

ABSTRACT:

The ABRF-2003ESRG sample is the 15th in a series of studies designed to allow participating members to evaluate their laboratory's abilities to analyze n-terminal proteins or peptide samples. This years study was done on a single protein that had been electroblotted to PVDF membrane. The original sample was submitted as part of a protein complex for analysis to an ESRG member's laboratory. One of the bands observed after SDS-PAGE analysis (another of which has already been used for the ABRF-2002ESRG study), running at an apparent molecular weight of 49kDa, was selected for this year's study. This sample demonstrated a single n-terminal sequence with initial yields around 2 pmol. Participants will be asked to assign sequence, identify the protein if possible, and return the tabular yield data to the ESRG for analysis. Comparison of initial yields, repetitive yields and overall performance of the instruments will be made, as well as run conditions and other factors that may affect the final results.

Materials and Methods

Sample Preparation:

This sample was prepared at the same time as the sample for the 2002 study. The protein complex* was present in several milliliters of solution. This was mixed with an equal volume of 2X Laemmli resolubilization buffer plus B-mercaptoethanol and boiled for 3 minutes. The protein was then run onto ten 15 lane precast 10-20% tris glycine 1.0 mm gels (NOVEX). After electrophoresis, the proteins were electrophoretically transferred to PVDF (NOVEX, 0.22 um) at 250 mAmps for 2 hours in a full immersion transfer tank system. PVDF membranes were then stained with 0.05% Coomassie blue G-250 in 50% methanol/10% acetic acid for five minutes, followed by destaining with 40% methanol/10% acetic acid and finally washing with water overnight. Bands used in this study were excised and placed into 0.5 ml microcentrifuge tubes and stored at room temperature for 1 year prior to being selected for this study. Individual bands (14 total) were sequenced for 25 amino acids by ESRG members on various models of sequencers to determine the expected performance of the sample. Amino acid analyses of three bands detected 9.4 pmole, 13.4 pmole, and 18.2 pmole of protein, with the average being 13.8 pmole. Assuming 15% sequencing yield from the PVDF, the expected average sequencing yield was approximately 2.0 pmole.

*Our thanks to Dr. Robert Macnab (Yale University, New Haven, CT, USA) and Dr. Tohru Minamino (Protonic Nanomachine ERATO, Kyoto, Japan) for the use of their protein complex in this study.

TABLE 1: ABRF-2003 ESRG Sequence Assignments from Facilities

| | | | | | | | | | | Expected Sequence | | | | | | | | | |